

Are rural places less safe for motorists? Definitions of urban and rural to understand road safety disparities

Carolyn McAndrews,¹ Kirsten Beyer,² Clare E Guse,³ Peter Layde⁴

¹Department of Urban and Regional Planning, University of Colorado Denver, Denver, Colorado, USA

²Medical College of Wisconsin Institute for Health and Society, Milwaukee, Wisconsin, USA

³Department of Family and Community Medicine, Injury Research Center, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

⁴Department of Emergency Medicine and Injury Research Center, Medical College of Wisconsin, Milwaukee, Wisconsin, USA

Correspondence to

Dr Carolyn McAndrews, Department of Urban and Regional Planning, University of Colorado Denver, 1250 14th Street, Suite 300, Denver, CO 80209, USA; carolyn.mcandrews@ucdenver.edu

Received 21 June 2016

Revised 20 October 2016

Accepted 30 December 2016

ABSTRACT

The objectives of the study are to understand road safety within the context of regional development processes and to assess how urban–rural categories represent differences in motor vehicle occupant fatality risk. We analysed 2015 motor vehicle occupant deaths in Wisconsin from 2010 to 2014, using three definitions of urban–rural continua and negative binomial regression to adjust for population density, travel exposure and the proportion of teen residents. Rural–Urban Commuting Area codes, Beale codes and the Census definition of urban and rural places do not explain differences in urban and rural transportation fatality rates when controlling for population density. Although it is widely believed that rural places are uniquely dangerous for motorised travel, this understanding may be an artefact of inaccurate constructs. Instead, population density is a more helpful way to represent transportation hazards across different types of settlement patterns, including commuter suburbs and exurbs.

INTRODUCTION

In the USA, in 2013, the fatality rate per 100 million vehicle-miles of travel (VMT) was 2.6 times higher in rural areas than in urban areas.¹ Explanations for this disparity focus on two aspects of transportation injury risk: (1) behavioural factors such as speeding, alcohol consumption and seat belt use and (2) environmental factors such as infrastructure design and lack of access to trauma centres.^{1–3}

Yet, previous research about rurality reminds us that rural places and populations are heterogeneous, that there is no agreed-on definition of rurality and that a simple urban–rural dichotomy, if invalid, prevents research and practice from addressing the underlying urban development factors that influence population health.^{4–6} For example, ‘rural’ places include traditional small towns that may be as safe as ‘urban’ places with similar built environments. Similarly, ‘urban’ places include zones of sparse development and natural landscapes that resemble rural places. To better capture this diversity of rurality, and to include the full spectrum of urban places (eg, commuter suburbs and exurbs), we used different definitions of urban–rural continua to explore whether using more specific categories would identify which types of rural places are uniquely hazardous for travel compared with their urban counterparts.

A key aspect of this analysis is that we adjusted for population density, in addition to using

urban–rural categories. Although population density is central to economic theory of urbanisation, many operational definitions of urban and rural places do not explicitly include it as a criterion. The US Census does include population density as a criterion, and sets 1000 persons per square mile as a starting point for inclusion in an urban area, but this definition does not reflect heterogeneity in population density within the category. Our framework allows us to account for hazards that associate with population density across the landscape such as high travel speeds, unsafe infrastructure design and higher travel distances (exposure), which we posit are relevant risk factors across the entire urban–rural continuum.^{7–10} Thus, this analysis brings together questions of sprawl and rurality into a general framework, which is necessary because the transportation injury risks associated with urbanisation patterns do not stop at the metropolitan edge.

METHODS

Our sample included all populated census tracts (N=1392) for the US state of Wisconsin. We classified each census tract according to two different definitions of urban–rural continua, each one capturing a specific aspect of rurality. The first continuum is based on Rural–Urban Commuting Area (RUCA) codes that reflect commuting patterns and economic relationships across places.¹¹ For instance, a place may be located outside of a large metropolitan area, but if a significant proportion of its population commutes to work inside the metropolitan area, this place is considered more urban than rural. The second continuum is based on a classification system designed by Chi and Marcouiller¹² to study urbanisation processes in Wisconsin. This classification system, which we refer to as the modified Beale code, describes urban and rural places as they would be perceived by urban migrants; it distinguishes between metropolitan and non-metropolitan areas, and further classifies each area by the size of its population centres. In addition to these two continua, we analysed differences in fatality rates using the Census definition of rural and urban areas. McAndrews *et al.*, 2016 provides further discussion of these operation definitions of urban-rural continua.¹³

We used a negative binomial model with a log-link function to estimate motor vehicle occupant fatality rates for each level of the urban–rural continuum, which was included as a categorical variable. We specified the model based on theoretical relationships, prior empirical research, bivariate

To cite: McAndrews C, Beyer K, Guse CE, *et al.* *Inj Prev* Published Online First: [please include Day Month Year] doi:10.1136/injuryprev-2016-042139

Brief report

relationships between motor vehicle occupant fatality rates and potential independent variables, and the Akaike information criterion (AIC) as a measure of model fit.

In addition to the log transformation of population density, we controlled for other independent variables that are hypothesised to influence motor vehicle occupant fatality rates. Traditionally, VMT is used to represent exposure, but population density already captures information about travel distances and is highly correlated with VMT (correlation coefficient (r) = -0.7 for our data). Therefore, we controlled for travel using the average daily number of trips per household in each census tract, which is less correlated with the distance of travel. We also controlled for the proportion of the population aged 16–19, representing teen drivers with higher injury risk. We initially considered median household income and the proportion of the population over age 65 as potential explanatory variables that could explain differences in urban and rural transportation fatality rates, but we did not include them in the final model because their associations with fatality rates were not statistically significant.

We used police-reported crash and fatality data for 2010–2014, collected at the state level and reported nationally with the Fatality Analysis Reporting System.¹⁴ We geocoded motor vehicle occupant fatalities based on the location of the crash. We manually assigned two fatalities to their nearest census tract, because their locations fell outside the census tract boundaries. We used 2010 census tract boundaries and population density

estimates from the US Census. Population (exposure) estimates are from the US Census for 2010, 2012, 2013 and 2014.¹⁵ We constructed the 2011 population estimate as the midpoint between 2010 and 2012 population for each tract. Estimates of average daily household motor vehicle trips are from the US Bureau of Transportation Statistics.¹⁶

RESULTS

Unadjusted fatality rates increased from about 10 per 100 000 population in the most urban zone to about 20 in the most rural zone (figure 1). This pattern was similar for the RUCA and modified Beale code definitions; the rural fatality rate was lower with the Census definition, 16 per 100 000 population.

After controlling for population density, household trips and proportion of teen population, rural census tracts had lower fatality rates compared with urban census tracts, but this was statistically significant for only levels 3 and 4 of the modified Beale code continuum. The rate ratios for these levels were 0.84 (95% CI 0.7 to 1.0) and 0.82 (95% CI 0.7 to 0.9), respectively (figure 1).

For the urban–rural continuum based on the RUCA code definition, census tracts with high levels of commuting that are located outside of core metro areas had a rate ratio 1.5 times (95% CI 1.2 to 2.0) the adjusted fatality rate of the most rural census tracts. These higher risk exurban tracts were coded as RUCA 2.0, 2.1, 3.0, 4.1, 4.1, 5.1, 5.2 and 6.1.




Figure 1 Unadjusted and regression-adjusted motor vehicle occupant fatality rates by urban–rural zone, Wisconsin, 2010–2014. RUCA, rural–urban commuting area.

The parameter estimates for population density and the proportion of teens in the population were significant in all three models ($p < 0.0001$ for all); population density had a negative association with motor vehicle occupant fatality rates, and the proportion of teens in the population had a positive association. Mean household daily trips had a negative association with fatality rates in all three models, but the parameter estimate was statistically significant only for the RUCA continuum ($p = 0.033$ for the RUCA; $p = 0.125$ for the modified Beale code; $p = 0.226$ for the Census definition). Travel behaviour in Wisconsin may not vary much across different built environment settings, and the relationship between household trips and motor vehicle occupant fatality rates may be stronger for regions with more diverse transportation systems.

DISCUSSION AND CONCLUSIONS

Although it is widely believed that 'rural' places are uniquely dangerous for motorised travel, this understanding may be an artefact of inaccurate constructs. Instead, when controlling for population density (and therefore indirectly controlling for risk factors such as speed and infrastructure design), rural places are similar to urban places, and may even be safer. Population density is a more helpful way to represent transportation hazards across different types of settlement patterns, and it does a better job including suburban and exurban developments that have received less attention from policy and practice. This dilemma with operational definitions of urban and rural places in the context of injury prevention is consistent with a prior study of adolescent smoking and drinking behaviour in which the choice of operational definitions of places influenced estimates of health outcomes and behaviours in meaningful ways.¹⁷

Operational definitions of urban–rural continua are important for transportation safety research and practice because they enable us to characterise what people and places have higher injury risk, and therefore they also enable the prioritisation of investment and intervention. For the specific case of injury prevention practice, professionals have started to revise the definitions of urban and rural to reflect practical needs. For instance, in Minnesota, emergency medical service providers customised the definition of urban and rural places to more accurately reflect the actual costs of providing service. Also, the state of Indiana revised the way it categorises the locations of where transportation crashes occur (urban, suburban, exurban and rural).^{18 19} The strength of this study is that we leveraged differences in the operational definitions of urban and rural continua to clarify their utility for transportation injury prevention across different types of settlement patterns. The results suggest that 'rural' places are not uniquely hazardous compared with more 'urban' places when accounting for settlement patterns—as represented by population density. The study area included the diverse urban–rural landscape of Wisconsin, but future work should test this framework for other geographical regions. In addition, future work should include direct measurement of the built environment and travel speeds to identify what facets of population density contribute to transportation injury risk.

Instead of understanding transportation hazards through the lens of the urban–rural dichotomy, injury prevention research, policy and programming can use population density, or similar objective measures, to understand and respond to specific hazards across a range of environments. This would enable road safety decision making to account for the consequences of regional development that influence the distribution of transportation injury risk.

What is already known on the subject?

- ▶ There is an inverse relationship between population density and motor vehicle occupant fatality rates.
- ▶ Motor vehicle occupant fatality rates for rural places in the USA are more than two times those of urban places.

What this study adds?

- ▶ Although it is widely believed that rural places are uniquely dangerous for motorised travel, this understanding may be an artefact of inaccurate constructs.
- ▶ Urban–rural differences in motor vehicle occupant fatality rates disappear when controlling for population density.
- ▶ Commuter suburbs and exurban places exhibit relatively high motor vehicle occupant fatality rates after adjusting for population density.
- ▶ Safety research and practice need updated constructs that better reflect true risk of existing and future settlement patterns.

Acknowledgements This research was supported by the Robert Wood Johnson Foundation Health and Society Scholars Program at the University of Wisconsin–Madison.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 National Highway Traffic Safety Administration. Rural/Urban Comparison. Traffic Safety Facts: 2013 Data. 2015; DOT HS 812 181. <http://www-nrd.nhtsa.dot.gov/Pubs/812181.pdf> (accessed 11 Jun 2016).
- 2 Watson Jr DC, Al-Kaisy A, Anderson ND. Examining the effect of speed, roadside features, and roadway geometry on crash experience along a rural corridor. *J Modern Transportation* 2014;22:84–95.
- 3 Siskind V, Steinhardt D, Sheehan M, et al. Risk factors for fatal crashes in rural Australia. *Accid Anal Prev* 2011;43:1082–8.
- 4 Hall SA, Kaufman JS, Ricketts TC. Defining urban and rural areas in US epidemiologic studies. *J Urban Health* 2006;83:162–75.
- 5 Hart LG, Larson EH, Lishner DM. Rural definitions for health policy and research. *Am J Public Health* 2005;95:1149–55.
- 6 Hartley D. Rural health disparities, population health, and rural culture. *Am J Public Health* 2004;94:1675–8.
- 7 Ewing R, Meakins G, Hamidi S, et al. Relationship between urban sprawl and physical activity, obesity, and morbidity—Update and refinement. *Health Place* 2014;26:118–26.
- 8 Ewing R, Schieber RA, Zegeer CV. Urban sprawl as a risk factor in motor vehicle occupant and pedestrian fatalities. *Am J Public Health* 2003;93:1541–5.
- 9 Ewing R, Dumbaugh E. The built environment and traffic safety: a review of empirical evidence. *J Plann Lit* 2009;23:347–67.
- 10 Fischer K, Sternfeld I, Melnick DS. Impact of population density on collision rates in a rapidly developing rural, exurban area of Los Angeles County. *Inj Prev* 2013;19:85–91.
- 11 WWAMI RUCA Rural Health Research Center. UW RHRC Rural Urban Commuting Area Codes. <http://depts.washington.edu/uwruca/> (accessed 11 Jun 2016).
- 12 Chi G, Marcoullier DW. Natural amenities and their effects on migration along the urban–rural continuum. *Ann Reg Sci* 2013;50:861–83.
- 13 McAndrews C, Layde P, Beyer K, et al. How do the definitions of urban and rural matter for transportation safety? Re-interpreting transportation fatalities as an outcome of regional development processes. *Accid Anal Prev* 2016;97:231–41.
- 14 National Highway Transportation Safety Administration. 2016. Fatality Analysis Reporting System. <http://www.nhtsa.gov/FARS> (accessed 20 May 2016).
- 15 Minnesota Population Center. *National Historical Geographic Information System: Version 2.0*. Minneapolis, MN: University of Minnesota 2011.

Brief report

- 16 Bureau of Transportation Statistics. Local Area Transportation Characteristics for Households (NHTS 2009 Transferability Statistics). 2016. http://www.rita.dot.gov/bts/sites/rita.dot.gov/bts/files/subject_areas/national_household_travel_survey/index.html (accessed 20 May 2016).
- 17 Brady JE, Weitzman BC. Inconsistencies in place definition: how different operational place definitions affect estimates of adolescent smoking and drinking risk. *Health Place* 2007, 13:562–8.
- 18 Rural Health Resource Center. An alternative approach to defining rural for the purpose of providing emergency medical services (EMS). Rural Road Environment Policy Paper. 2004. <http://www.rospa.com/rospaweb/docs/advice-services/road-safety/rural-road-environment.pdf>
- 19 Newby B. Improving the locational classification of Indiana motor vehicle collisions. 2011. Indiana 2010 Traffic Safety Facts, 11-C10. Indiana Criminal Justice Institute, Indiana University Center for Criminal Justice Research. <http://policyinstitute.iu.edu/publication/details/553>



Are rural places less safe for motorists? Definitions of urban and rural to understand road safety disparities

Carolyn McAndrews, Kirsten Beyer, Clare E Guse and Peter Layde

Inj Prev published online January 24, 2017

Updated information and services can be found at:
<http://injuryprevention.bmj.com/content/early/2017/01/24/injuryprev-2016-042139>

These include:

References

This article cites 12 articles, 1 of which you can access for free at:
<http://injuryprevention.bmj.com/content/early/2017/01/24/injuryprev-2016-042139#BIBL>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Notes

To request permissions go to:
<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:
<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:
<http://group.bmj.com/subscribe/>